blank page

POISON KEEP OUT OF REACH OF CHILDREN READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Apparent

C. RECEIPTION

Affray 300

HERBICIDE

ACTIVE CONSTITUENT: 300 g/L 2,4-D SOLVENT: 259 g/L HYDROCARBON LIQUID

GROUP

HERBICIDE

For the integrated control of Groundsel Bush, Mother-of-Millions, Noogoora Burr, Bathurst Burr, Water Hyacinth and other weeds as listed in the Directions for Use table.

This is a PHENOXY HERBICIDE that can cause severe damage to susceptible crops such as cotton, grapes, tomatoes, oilseed crops and ornamentals.

IMPORTANT: Read this booklet before use.

MA Approval No: 56510/108336

APPARENT PTY LTD A.C.N. 143 724 136 Suite G.08 | 762 Toorak Road, Glen Iris VIC 3146 Australia M 0411 227 338 | E enquiries@apparentag.com.au www.apparentag.com.au

DIRECTIONS FOR USE

RESTRAINT: Avoid use in sown pasture.

Situation	Weeds Controlled	State	Rate	Critical Comments
Non-crop & Pasture	Groundsel Bush	Qld & NSW only	10 L/ha	Helicopter application. Raindrop D8 nozzles angles back at 45° to minimise spray drift. Apply in no less
				than 100 L/ha.
			100 mL/10 L water	Knapsack spray. Thorough even coverage of the plant is necessary.
			1 L/10 L water	Sprinkler sprayer. Coverage: 1 L spray solution/100 m ² .
			33 mL/L kerosene or	Basal bark spray: Apply the mixture around the base of the stem to a height of 25 cm above the ground.
			mineral turpentine	Make sure all the stem to this height is well wetted.
				Cut stump: Cut the stem as close to the ground as possible and thoroughly wet the fresh cut stem with
				the mixture.
	Mother-of-Millions		70 mL/10 L water	Knapsack spray. Thorough even coverage of leaves and plantlets is necessary.
			7 L/1000 L water/ha	Hand Gun. Thorough even coverage of leaves and plantlets is necessary.
	Noogoora Burr, Bathurst Burr		3.3 L/ha	Helicopter application or sprinkler sprayer.
	Castor Oil Plant	Qld only	1 L/10 L kerosene	Apply as a basal bark treatment.
	Galvanised Burr, Chinese Celtis (Elm)		1 L/100 L water	Apply as an overall spray on young plants when actively growing.
	seedlings			
	Wild Tobacco Tree		1 L/40 L kerosene	Apply as a cut stump or basal bark when plant actively growing.
	Balloon Cotton, Datura (seedlings),		7 L/1000 L water	Apply as an overall spray when plants actively growing.
	Sida Retusa, Creeping Lantana,			
	Ageratum spp., Fireweed			
	Saffron Thistle		1 L/100 L water	Apply as an overall spray when plants actively growing.
	Water Hyacinth, Water Lettuce	NSW, Qld, WA	50 mL/10 L water	Knapsack spray. Coverage: 10 L spray solution/100 m ² . Avoid causing submersion of sprayed plants.
water, drains, dams		and NT only	1 L/200 L water	Hand Gun. Coverage: 200 L spray solution/1000 m ² . Avoid causing submersion of sprayed plants.
margins, lakes &			1 L/20 L water	Sprinkler sprayer. Coverage: 20 L spray solution/1000 m ² .
streams			5 L/200 L water/ha	Helicopter application. Raindrop D8 nozzles angled back at 45° to minimise spray drift.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIOD:

DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 7 DAYS AFTER APPLICATION.

GENERAL INSTRUCTIONS:

Before opening, carefully read Directions for Use, Precautionary Statements, Safety Directions and First Aid Instructions.

EQUIPMENT MAINTENANCE AND USE

Equipment that has been used for this chemical should not be used for the application of other materials to sensitive plants, unless it has been well washed out with hot soapy water or 1% solution of ammonia followed by several clear water rinses.

COMPATIBILITY

Compatible with biological control agents and can be used in Integrated Pest Management Programs (IPM).

RESISTANT WEEDS WARNING

GROUP HERBICIDE

Apparent Affray 300 Herbicide is a member of the phenoxys group of herbicides. Apparent Affray 300 Herbicide has the disruptors of plant cell growth mode of action. For weed resistance management, Apparent Affray 300 Herbicide is a Group I Herbicide.

Some naturally occurring weed biotypes resistant to Apparent Affray 300 Herbicide and other Group I Herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by Apparent Affray 300 Herbicide or other Group I Herbicides.

Since the occurrence of resistant weeds is difficult to detect prior to use, Apparent Pty Ltd accepts no liability for any losses that may result from the failure of Apparent Affray 300 Herbicide to control resistant weeds.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DO NOT spray in high winds. DO NOT spray cereals if lucerne is present. DO NOT spray crops or weeds outside the stages indicated in "Critical Comments" as damage, loss of yield or inadequate weed control may result.

Drift Warning: DO NOT apply under weather conditions, or from spraying equipment that may cause spray to drift onto nearby susceptible plants/crops, cropping lands or pastures. Avoid spray drift onto susceptible crops such as cotton, tobacco, tomatoes, vines, fruit trees, vegetables, legume crops and pastures, oilseed crops and susceptible trees (e.g. Kurrajongs, Belahs and Eucalypts).

Avoid use on Sown Pastures.

PROTECTION OF LIVESTOCK

Low hazard to bees. May be applied at any time as recommended in the Directions for Use.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate streams, rivers, or waterways with the chemical or used container.

Drift Warning: D0 NOT use unless wind speed is more than 3 kilometres per hour and less than 15 kilometres per hour as measured at the application site.

DO NOT apply with smaller than coarse to very coarse spray droplets according to the ASAE S572 definition for standard nozzles.

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area away from children, animals, food, feedstuffs, seed and fertilisers. DO NOT store for prolonged periods in direct sunlight.

Triple-rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point.

If not recycling, break, crush, or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.

SAFETY DIRECTIONS

Poisonous if swallowed. Avoid contact with eyes and skin. DO NOT inhale spray mist. When preparing spray wear PVC or rubber apron, elbow-length PVC gloves and face shield. When using the prepared spray wear face shield. If product on skin, immediately wash area with soap and water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves, face shield and contaminated clothing.

FIRST AID

If poisoning occurs, contact a Doctor or Poisons Information Centre. Phone Australia 13 11 26.

SAFETY DATA SHEET

Additional information is listed in the Safety Data Sheet (SDS) available from the supplier.

CONDITIONS OF SALE

The use of Apparent Affray 300 Herbicide being beyond the control of the manufacturer no warranty expressed or implied is given by Apparent Pty Ltd regarding its suitability, fitness or efficiency for any purpose for which it is used by the buyer, whether in accordance with the directions or not and Apparent Pty Ltd accepts with no responsibility for any consequences whatsoever resulting from the use of this product.